


A **stochastic timed automaton** $(\mathcal{E}, \mathcal{X}, \Gamma, P, p_{x_0}, F)$ transforms the stochastic clock structure F into the pmf's of the event and state sequences:



The basic brick to carry out this task, is the computation of probabilities of the type:

$$P(E_k = e | X_{k-1} = x)$$

Indeed, if these probabilities are known, then, using the total probability rule:

$$\bullet P(E_k = e) = \sum_{x \in \mathcal{X}} P(E_k = e | X_{k-1} = x) P(X_{k-1} = x)$$

$$\bullet P(X_k = x') = \sum_{x \in \mathcal{X}} P(X_k = x' | X_{k-1} = x) P(X_{k-1} = x)$$

$$P(X_k = x' | X_{k-1} = x) = \sum_{e \in \Gamma(x)} \underbrace{P(X_k = x' | X_{k-1} = x, E_k = e)}_{p(x' | x, e)} P(E_k = e | X_{k-1} = x)$$

$\Rightarrow$ Given p_{x_0} , we compute p_{E_1} and p_{x_1} .

Then, given p_{x_1} , we compute p_{E_2} and p_{x_2} .

And so on... $\Rightarrow$ iterative approach

Formally,

$$P(E_k=e | X_{k-1}=x) = P\left(Y_{e,k-1} < \min_{\substack{e' \in \Gamma(x) \\ e' \neq e}} Y_{e',k-1}\right)$$

Unfortunately, the distributions of the residual lifetimes are unknown in general, if we only assume the knowledge of the current state. Hence, $P(E_k=e | X_{k-1}=x)$ cannot be computed easily.

For this reason, to circumvent the problem, in the exercises we had to consider all the possible cases, starting from initialization, which results into a dramatic proliferation of cases as the event index k increases.

The question now is whether there exist classes of stochastic timed automata for which computing $P(E_k=e | X_{k-1}=x)$ is an easy task.

The answer is given by stochastic timed automata with Poisson clock structure.

DEFINITION - A Poisson clock structure is a stochastic clock structure $F = \{F_e : e \in \mathcal{E}\}$ where all the distributions F_e are exponential, i.e. for all $e \in \mathcal{E}$:

$$F_e(t) = \begin{cases} 1 - e^{-\lambda_e t} & \text{if } t \geq 0 \\ 0 & \text{otherwise.} \end{cases} \quad \lambda_e > 0$$

MAIN RESULT

For a stochastic timed automaton with Poisson clock structure, the residual lifetimes of an event have the same distribution as the corresponding total lifetimes.

In other words: $V_e \sim \text{Exp}\left(\frac{1}{\lambda_e}\right) \Rightarrow Y_e \sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$

Sketch of the proof (by induction)

1) At initialization ($k=0$):

$$Y_{e,0} = V_{e,1} \text{ for all } e \in \Gamma(X_0)$$

Since $V_{e,1} \sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$, then $Y_{e,0} \sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$

$\Rightarrow$ The statement is true for $k=0$.


2) We assume that the statement is true for $k-1$, and we prove that it is true for k .

Case i) Event e is activated in state X_k .

$$\Rightarrow Y_{e,k} = V_{e,N_{e,k}}$$

Since $V_{e,N_{e,k}} \sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$, then $Y_{e,k} \sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$

Case ii) Event e "continues" in state X_k .



$$\Rightarrow Y_{e,k} = Y_{e,k-1} - Y_{k-1}^*$$

$$P(Y_{e,k} \leq t \mid Y_{e,k-1} > Y_{k-1}^*) =$$

$$= P(Y_{e,k-1} - Y_{k-1}^* \leq t \mid Y_{e,k-1} > Y_{k-1}^*)$$

$$\xrightarrow{\text{Extended memoryless property}} = P(Y_{e,k-1} \leq t) = 1 - e^{-\lambda_e t}, \quad t \geq 0$$

Extended
memoryless
property

By induction

$$\text{Hence, } Y_{e,k} \sim \text{Exp}\left(\frac{1}{\lambda_e}\right).$$

$\Rightarrow$ The statement is true for k .

As a consequence of the main result, the following holds:

$$P(E_k = e \mid X_{k-1} = x) = \frac{\lambda_e}{\mathcal{L}(x)}$$

where $\mathcal{L}(x) \triangleq \sum_{e' \in \Gamma(x)} \lambda_{e'}$

$\rightarrow$ We have a closed-form formula for $P(E_k = e \mid X_{k-1} = x)$!

Proof

$$P(E_k=e | X_{k-1}=x)$$

$$= P\left(Y_{e,k-1} < \min_{\substack{e' \in \Gamma(x) \\ e' \neq e}} Y_{e',k-1} \right)$$

$\sim \text{Exp}\left(\frac{1}{\lambda_e}\right)$
 $\sim \text{Exp}\left(\frac{1}{\lambda_{e'}}\right)$

Superposition
 $\Rightarrow \text{exp. distr. with rate } \sum_{\substack{e' \in \Gamma(x) \\ e' \neq e}} \lambda_{e'}$

$$= \frac{\lambda_e}{\lambda_e + \sum_{\substack{e' \in \Gamma(x) \\ e' \neq e}} \lambda_{e'}} = \frac{\lambda_e}{\Lambda(x)}$$

$P(X < Y) = \frac{\lambda}{\lambda + \mu}$

with $X \sim \text{Exp}\left(\frac{1}{\lambda}\right)$, $Y \sim \text{Exp}\left(\frac{1}{\mu}\right)$, independent

Exploiting the formula for $P(E_k=e | X_{k-1}=x)$ derived above, we can apply the iterative approach illustrated at the beginning for the computation of p_{E_k} and p_{X_k} .

Let $\mathcal{E} = \{1, 2, \dots, m\}$ and $\mathcal{X} = \{1, 2, \dots, n\}$, and define the row vectors:

$$\pi_E(k) \triangleq [P(E_k=1) \ P(E_k=2) \ \dots \ P(E_k=m)] \in \mathbb{R}^{1 \times m}$$

$$\pi_X(k) \triangleq [P(X_k=1) \ P(X_k=2) \ \dots \ P(X_k=n)] \in \mathbb{R}^{1 \times n}$$

Moreover, define the matrices:

$$P_E \triangleq \begin{bmatrix} \frac{\lambda_1}{\Lambda(1)} & \frac{\lambda_2}{\Lambda(1)} & \dots & \frac{\lambda_m}{\Lambda(1)} \\ \frac{\lambda_1}{\Lambda(2)} & \frac{\lambda_2}{\Lambda(2)} & \dots & \frac{\lambda_m}{\Lambda(2)} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\lambda_1}{\Lambda(n)} & \frac{\lambda_2}{\Lambda(n)} & \dots & \frac{\lambda_m}{\Lambda(n)} \end{bmatrix} \in \mathbb{R}^{n \times m}$$

and

$$P_X \triangleq \begin{bmatrix} p_{1,1} & p_{1,2} & \dots & p_{1,n} \\ p_{2,1} & p_{2,2} & \dots & p_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ p_{n,1} & p_{n,2} & \dots & p_{n,n} \end{bmatrix} \in \mathbb{R}^{n \times n}$$

$$\text{where } p_{i,j} \triangleq \sum_{e \in \Gamma(i)} p(j|i, e) \cdot \frac{\lambda_e}{\Lambda(i)}$$

Then we have:

$$P(E_k = j) = \sum_{i=1}^n \underbrace{P(E_k = j | X_{k-1} = i)}_{\substack{\frac{\lambda_j}{\Lambda(i)} \\ \text{entry in position} \\ (i, j) \text{ of } P_E}} P(X_{k-1} = i)$$

$\swarrow$ j -th entry of $\pi_E(k)$

$\searrow$ i -th entry of $\pi_X(k-1)$

$\swarrow$ product of $\pi_X(k-1)$ and the j -th column of P_E

=> In matrix form:

$$\pi_E(k) = \pi_X(k-1) P_E$$

Similarly:

$$P(X_k=j) = \sum_{i=1}^n \underbrace{P(X_k=j | X_{k-1}=i)}_{\substack{P_{i,j} \\ \text{entry in position} \\ (i,j) \text{ of } P_X}} \underbrace{P(X_{k-1}=i)}_{\substack{i\text{-th entry} \\ \text{of } \pi_X(k-1)}}$$

$P(X_k=j)$ is the j -th entry of $\pi_X(k)$.

The entire sum is the product of $\pi_X(k-1)$ and the j -th column of P_X .

=> In matrix form:

$$\pi_X(k) = \pi_X(k-1) P_X$$

Notice that:

$$\begin{aligned} \pi_X(0) &= [P(X_0=1) \ P(X_0=2) \ \dots \ P(X_0=n)] \\ &= [p_{X_0}(1) \ p_{X_0}(2) \ \dots \ p_{X_0}(n)] \Rightarrow \text{known} \end{aligned}$$

Hence:

$$\pi_X(1) = \pi_X(0) P_X$$

$$\pi_X(2) = \pi_X(1) P_X = \pi_X(0) P_X \cdot P_X = \pi_X(0) P_X^2$$

$$\pi_X(3) = \pi_X(2) P_X = \pi_X(0) P_X^2 \cdot P_X = \pi_X(0) P_X^3$$

and so on.

We easily figure out that:

$$\pi_x(k) = \pi_x(0) P_x^k \quad k=1,2,3,\dots$$

Moreover, from $\pi_E(k) = \pi_x(k-1) P_E$:

$$\pi_E(k) = \pi_x(0) P_x^{k-1} \cdot P_E \quad k=1,2,3,\dots$$


$\Rightarrow$ Computing p_{E_k} and p_{x_k} boils down to a problem of linear algebra

$\Rightarrow$ only matrix computations

$\Rightarrow$ computationally very efficient!

STATE HOLDING TIME

DEFINITION - The **state holding time** $V(x)$ is the time that the system remains in state $x \in \mathcal{X}$.



We want to compute the **distribution of $V(x)$** for a stochastic timed automaton with Poisson clock structure.


PRELIMINARY RESULT

Let $X_1, X_2, \dots, X_n, X_{n+1}$ be independent, identically distributed random variables with $X_i \sim \text{Exp}\left(\frac{1}{\lambda}\right)$ for all $i=1, 2, \dots, n+1$.

Then, for $t > 0$:

$$P(X_1 + \dots + X_n < t, X_1 + \dots + X_n + X_{n+1} > t) = \frac{(\lambda t)^n}{n!} e^{-\lambda t}$$

Recalling that $P(V(x) \leq t) = 1 - P(V(x) > t)$, we start from computing $P(V(x) > t)$, for $t > 0$.



$$P(V(x) > t) =$$

$$= P(\text{no state transition}^{(*)} \text{ occurs over } (0, t])$$


(*) we mean no transition to a state $x'' \neq x$

$$= P\left(\bigcap_{e \in \Gamma(x)} \left\{ \text{no state transition triggered by event } e \text{ over } (0, t] \right\}\right)$$

$$= \prod_{e \in \Gamma(x)} P(\text{no state transition triggered by event } e^{(\Delta)} \text{ over } (0, t])$$

independent

(Δ)




$$= \prod_{e \in \Gamma(x)} P\left(\bigcup_{n=0}^{\infty} \left\{ \text{event } e \text{ occurs exactly } n \text{ times over } (0, t], \right. \right. \\ \left. \left. \text{it does not trigger a state transition} \right\}\right)$$

disjoint

$$= \prod_{e \in \Gamma(x)} \sum_{n=0}^{\infty} P\left(\text{event } e \text{ occurs exactly } n \text{ times over } (0, t], \right. \\ \left. \text{it does not trigger a state transition} \right)$$

(□)



where $e^{(i)}$ = i -th occurrence of event e , and $Y_{e^{(i)}}$ its lifetime

$$\Rightarrow P(\text{event } e \text{ occurs exactly } n \text{ times over } (0, t]) \\ = P(Y_{e^{(1)}} + \dots + Y_{e^{(n)}} < t, Y_{e^{(1)}} + \dots + Y_{e^{(n)}} + Y_{e^{(n+1)}} > t) \\ = \frac{(\lambda t)^n}{n!} e^{-\lambda t}$$

$$= \prod_{e \in \Gamma(x)} \sum_{n=0}^{\infty} \underbrace{\frac{(\lambda t)^n}{n!} e^{-\lambda t}}_{\text{prob. event } e \text{ occurs } n \text{ times over } (0, t]} \cdot \underbrace{p(x|x, e)}_{\text{prob. to remain in state } x \text{ when event } e \text{ occurs}}$$

(Bernoulli scheme)

n times

$$= \prod_{e \in \Gamma(x)} e^{-\lambda_e t} \sum_{n=0}^{\infty} \frac{[\overset{(o)}{\lambda_e p(x|x,e)} t]^n}{n!}$$

(o)

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$= \prod_{e \in \Gamma(x)} e^{-\lambda_e t} \cdot e^{\lambda_e p(x|x,e) t}$$

$$= \prod_{e \in \Gamma(x)} e^{-\lambda_e [1-p(x|x,e)] t}$$

$$= e^{-\left\{ \sum_{e \in \Gamma(x)} \lambda_e [1-p(x|x,e)] \right\} t}$$

$$\Rightarrow P(V(x) \leq t) = 1 - e^{-\left\{ \sum_{e \in \Gamma(x)} \lambda_e [1-p(x|x,e)] \right\} t}, \quad t \geq 0$$

We have thus shown that $V(x)$ has an exponential distribution with rate $\sum_{e \in \Gamma(x)} \lambda_e [1-p(x|x,e)]$


Notice that:

$$\sum_{x' \in \mathcal{X}} p(x'|x,e) = 1 \Rightarrow p(x|x,e) + \sum_{x' \neq x} p(x'|x,e) = 1$$

$$\Rightarrow 1 - p(x|x,e) = \sum_{x' \neq x} p(x'|x,e)$$

$$\Rightarrow \sum_{e \in \Gamma(x)} \lambda_e [1-p(x|x,e)] = \sum_{e \in \Gamma(x)} \sum_{x' \neq x} \lambda_e p(x'|x,e)$$

EXAMPLE



event	rate
a	λ_a
b	λ_b
c	λ_c

The rate of the exponential distribution of $V(1)$ is:

$$\lambda_a(1-p) + \lambda_b \Rightarrow E[V(1)] = \frac{1}{\lambda_a(1-p) + \lambda_b}$$

$\uparrow$
 expected value

For $V(2)$:

$$\lambda_a + \lambda_c \Rightarrow E[V(2)] = \frac{1}{\lambda_a + \lambda_c}$$

For $V(3)$:

$$\lambda_a + \lambda_b(1-q) + \lambda_c \Rightarrow E[V(3)] = \frac{1}{\lambda_a + \lambda_b(1-q) + \lambda_c}$$

POISSON PROCESSES

A **Poisson process** is the counting process of an event which is always possible, and characterized by interevent times that are:

- independent
- identically distributed with an exponential distribution.


The rate of the exponential distribution is also called the rate of the Poisson process.

A Poisson process can be viewed as a stochastic timed automaton with Poisson clock structure:

$$(\mathcal{E}, \mathcal{X}, \Gamma, f, x_0, F)$$

where $\mathcal{E} = \{e\}$, $\mathcal{X} = \{0, 1, 2, 3, \dots\}$, the state transition diagram is given by


the state counts the number of occurrences of event e



and $F = \{F_e\}$ with $F_e(t) = 1 - e^{-\lambda t}$, $t \geq 0$, $\lambda > 0$.

Let

$N_e(t, t+T) \triangleq$ # occurrences of event e over the interval $(t, t+T]$



$N_e(t, t+T)$ is a discrete random variable taking values in $\{0, 1, 2, \dots\}$. It can be shown that the pmf of $N_e(t, t+T)$ has the following form:

$$P[N_e(t, t+T) = n] = \frac{(\lambda T)^n}{n!} e^{-\lambda T}, \quad n = 0, 1, 2, \dots$$

→ Poisson distribution with parameter λT


The expected value of $N_e(t, t+T)$ is

$$E[N_e(t, t+T)] = \lambda T.$$

Notice that the above formula does not depend on the time instant t , but only on the length T of the interval.

$\Rightarrow$ It is a consequence of the memoryless property of the exponential distribution


- Consider, for example, $P(N_e(0, T) = 0)$.



$$\Rightarrow P(N_e(0,T)=0) = P(V_{e,1} > T) = 1 - P(V_{e,1} \leq T) \\ = 1 - (1 - e^{-\lambda T}) = e^{-\lambda T}$$

- Now consider $P(N_e(t,t+T)=0)$ for $t > 0$.

Say we had k events e before time t (k is unspecified).



$$\Rightarrow P(N_e(t,t+T)=0) =$$

$$= P\left(\sum_{h=1}^k V_{e,h} + V_{e,k+1} > t+T \mid \sum_{h=1}^k V_{e,h} + V_{e,k+1} > t\right)$$

$$= P\left[V_{e,k+1} > \left(t - \sum_{h=1}^k V_{e,h}\right) + T \mid V_{e,k+1} > t - \sum_{h=1}^k V_{e,h}\right]$$

$$= P(V_{e,k+1} > T) = 1 - P(V_{e,k+1} \leq T)$$

$$= 1 - (1 - e^{-\lambda T}) = e^{-\lambda T}$$

extended
memoryless
property


This random variable
is positive, because
it is conditional to the
fact that k events e
occurred before t

Hence, $P(N_e(t,t+T)=0) = e^{-\lambda T}$ independent of $t \geq 0$.

In the following, we will omit t in the notation of N_e :

$$N_e(t, t+T) \longrightarrow N_e(T) \text{ independent of } t.$$

Notice that $P(N_e(T)=n)$ corresponds to the following sample path:



$$\Rightarrow \frac{(\lambda T)^n}{n!} = P(N_e(T)=n) = P\left(\sum_{i=1}^n V_{e,i} < T, \sum_{i=1}^{n+1} V_{e,i} > T\right)$$

We have thus proven the preliminary result on top of page 9. Simply pose $V_{e,i} \leftarrow X_i$, $i=1, \dots, n+1$, and $T \leftarrow t$.

BYPRODUCT

Let $X_1, X_2, \dots, X_n, X_{n+1}$ be independent, identically distributed random variables with $X_i \sim \text{Exp}\left(\frac{1}{\lambda}\right)$ for all $i=1, 2, \dots, n+1$.

Then, for $t > 0$:


$$P(X_1 + \dots + X_n < t, X_1 + \dots + X_n + X_{n+1} > t) = \frac{(\lambda t)^n}{n!} e^{-\lambda t}$$

EXAMPLE (computations with Poisson processes)

Assume that event e is generated by a Poisson process with rate $\lambda > 0$.

- Compute the probability that event e does not occur over $(0, T]$, and occurs exactly once over $(T, 2T]$.

This probability corresponds to the following sample path:




$$\begin{aligned} \Rightarrow P(\dots) &= \begin{cases} P(T < V_{e,1} < 2T, V_{e,1} + V_{e,2} > 2T) \\ \text{using Poisson:} \end{cases} \\ &= P(N_e(0, T) = 0) P(N_e(T, 2T) = 1) \\ &= P(N_e(T) = 0) P(N_e(T) = 1) = e^{-\lambda T} \cdot (\lambda T) e^{-\lambda T} \end{aligned}$$

the two intervals are treated independently

- Compute the probability that event e occurs at least twice over $(0, T]$.

This probability corresponds to the following sample path:



Notice that, once we impose that the second event occurs before T , then the condition is satisfied, and we don't care about what happens next.

$$\Rightarrow P(\dots) = \begin{cases} P(V_{e,1} + V_{e,2} < T) \\ \text{using Poisson:} \end{cases}$$

$$\begin{aligned} P(N_e(T) \geq 2) &= 1 - P(N_e(T) = 0) - P(N_e(T) = 1) \\ &= 1 - e^{-\lambda T} - (\lambda T) e^{-\lambda T} \\ &= 1 - e^{-\lambda T} (1 + \lambda T) \end{aligned}$$

BYPRODUCT

Let X_1, X_2 be independent, identically distributed random variables with $X_i \sim \text{Exp}\left(\frac{1}{\lambda}\right)$, $i=1,2$. Then, for $t > 0$:

$$P(X_1 + X_2 < t) = 1 - e^{-\lambda t} (1 + \lambda t)$$

Notice that, if the question was:

"Compute the probability that event e occurs exactly twice over $(0, T]$ "

then the answer was:


$$P(\dots) = \begin{cases} P(V_{e,1} + V_{e,2} < T, V_{e,1} + V_{e,2} + V_{e,3} > T) \\ \text{using Poisson: } P(N_e(T) = 2) = \frac{(\lambda T)^2}{2} e^{-\lambda T} \end{cases}$$

USEFUL RESULT

Assume that event e is generated by a Poisson process with rate $\lambda > 0$.

Moreover, assume that, after its occurrence, event e is classified of type 1 with probability q , and of type 2 with probability $1-q$.

Then, the Poisson process generating event e can be viewed as the **superposition** of two independent Poisson processes, one with rate λq (generating events e classified of type 1), and the other with rate $\lambda(1-q)$ (generating events e classified of type 2).



=> Lifetime of event e : $V_e = \min \{ V_{e,1}, V_{e,2} \}$

superposition



exp. distributed
with rate

$$\lambda q + \lambda(1-q) = \lambda \quad \checkmark$$

exp. distributed
with rate λq

exp. distributed
with rate $\lambda(1-q)$

independent