


• Serie di Laurent

→ serie di Taylor per numeri complessi.

LAURENT SERIES



Laurent Series Explained | How to Determine Laurent Series | C...

Everything you need to know about Laurent Series explained. The video will contain pr...

m.youtube.com

→ The most complete example:

Example: Determine all possible Laurent Series for each function.

3) $f(z) = \frac{1}{(z-1)(z-2)}$, around $z=0$



for $|z| < 1$:

$$\frac{1}{z-2} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \text{ for } |z| < 2$$

$$\frac{1}{z-1} = -1 \cdot \frac{1}{1-z} = \left\{ (1^*) \right\} = -\sum_{n=0}^{\infty} z^n, \text{ for } |z| < 1$$

for $1 < |z| < 2$:

$$\frac{1}{z-2} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \text{ for } |z| < 2$$

$$\frac{1}{z-1} = \frac{1}{z} \cdot \frac{1}{1-\frac{1}{z}} = \left\{ (2^*) \right\} = \frac{1}{z} \cdot \sum_{n=0}^{\infty} \left(\frac{1}{z}\right)^n = \sum_{n=1}^{\infty} \frac{1}{z^n}, \text{ for } |z| > 1$$

$$f(z) = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} - \left(-\sum_{n=0}^{\infty} z^n \right) = \sum_{n=0}^{\infty} \left(1 - \frac{1}{2^{n+1}} \right) \cdot z^n, \text{ for } |z| < 1$$

$$f(z) = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} - \sum_{n=1}^{\infty} \frac{1}{z^n}, \text{ for } 1 < |z| < 2$$

$$\frac{1}{1-c \cdot w} = \sum_{n=0}^{\infty} (c \cdot w)^n, \text{ for } |w| < \frac{1}{|c|} \quad (1^*)$$

$$\frac{1}{1-\frac{b}{w}} = \sum_{n=0}^{\infty} \left(\frac{b}{w}\right)^n, \text{ for } |w| > |b| \quad (2^*)$$

note: $\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}$
Partial fraction expansion

- Wak-up:

- Definition of Laurent Series:

$$f(z) = \dots + \frac{a_{-3}}{(z-z_0)^3} + \frac{a_{-2}}{(z-z_0)^2} + \frac{a_{-1}}{(z-z_0)^1} + a_0 + a_1 \cdot (z-z_0)^1 + a_2 \cdot (z-z_0)^2 + a_3 \cdot (z-z_0)^3 + \dots$$

Laurent Series


- Such that :

Theorem: Laurent Series

if $f(z)$ is analytic on $R_1 < |z-z_0| < R_2$, then

$$f(z) = \underbrace{\sum_{n=0}^{\infty} a_n \cdot (z-z_0)^n}_{\text{analytic part}} + \underbrace{\sum_{n=1}^{\infty} a_{-n} \cdot (z-z_0)^{-n}}_{\text{principal part}}, \text{ for all } z \text{ on } R_1 < |z-z_0| < R_2$$

and where $a_n = \frac{1}{2\pi i} \oint_C \frac{f(s)}{(s-z_0)^{n+1}} ds \quad (n=0, \pm 1, \pm 2, \dots)$.



- Usually we don't use the formula for calculating a_n

- We can more easily use the harmonic series:

$$\frac{1}{1-w} = w^0 + w^1 + w^2 + \dots = \sum_{n=0}^{\infty} (w)^n, \text{ for } |w| < 1 \quad (1)$$

$$\frac{1}{1-\frac{1}{w}} = \frac{1}{w^0} + \frac{1}{w^1} + \frac{1}{w^2} + \dots = \sum_{n=0}^{\infty} \left(\frac{1}{w}\right)^n, \text{ for } |w| > 1 \quad (2)$$


- So we can solve the following problem:

Example: Determine all possible Laurent Series for each function.

$$\frac{1}{1-c \cdot w} = \sum_{n=0}^{\infty} (c \cdot w)^n, \text{ for } |w| < \frac{1}{|c|} \quad (1^*)$$

$$\frac{1}{1-\frac{b}{w}} = \sum_{n=0}^{\infty} \left(\frac{b}{w}\right)^n, \text{ for } |w| > |b| \quad (2^*)$$

1) $f(z) = \frac{1}{z-2}$, around $z=0$




for $|z| < 2$:

$$f(z) = \frac{1}{z-2} = -1 \cdot \frac{1}{2-z} = -\frac{1}{2} \cdot \frac{1}{1-\frac{z}{2}} = \left\{ (1^*), w = \frac{z}{2}, c = \frac{1}{2} \right\} = -\frac{1}{2} \cdot \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \text{ for } |z| < 2$$

for $|z| > 2$:

$$f(z) = \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\frac{2}{z}} = \left\{ (2^*), w = \frac{2}{z}, b = 2 \right\} = \frac{1}{z} \cdot \sum_{n=0}^{\infty} \left(\frac{2}{z}\right)^n = \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \left\{ k=n+1 \right\} = \sum_{n=1}^{\infty} \frac{2^{n-1}}{z^n}, \text{ for } |z| > 2$$

2) $f(z) = \frac{1}{z-2}$, around $z=1$



for $|z-1| < 1$:

$$f(z) = \frac{1}{z-2} = \frac{1}{(z-1)-1} = -1 \cdot \frac{1}{1-(z-1)} = \left\{ (1^k) \right\} = -\sum_{n=0}^{\infty} (z-1)^n, \text{ for } |z-1| < 1$$

for $|z-1| > 1$:

$$\begin{aligned} f(z) &= \frac{1}{z-2} = \frac{1}{(z-1)-1} = \frac{1}{z-1} \cdot \frac{1}{1-\frac{1}{z-1}} = \left\{ (2^k) \right\} = \frac{1}{z-1} \cdot \sum_{n=0}^{\infty} \left(\frac{1}{z-1} \right)^n \\ &= \sum_{n=0}^{\infty} \frac{1}{(z-1)^{n+1}} = \sum_{n=1}^{\infty} \frac{1}{(z-1)^n} \end{aligned}$$

• Or another problem as:

Example: Determine all possible Laurent Series for each function.

3) $f(z) = \frac{1}{(z-1)(z-2)}$, around $z=0$



for $|z| < 1$:

$$\frac{1}{z-2} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \text{ for } |z| < 2$$

$$\frac{1}{z-1} = -1 \cdot \frac{1}{1-z} = \left\{ (1^k) \right\} = -\sum_{n=0}^{\infty} z^n, \text{ for } |z| < 1$$

$$f(z) = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} - \left(-\sum_{n=0}^{\infty} z^n \right) = \sum_{n=0}^{\infty} \left(1 - \frac{1}{2^{n+1}} \right) \cdot z^n, \text{ for } |z| < 1$$

for $1 < |z| < 2$:

$$\frac{1}{z-2} = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}, \text{ for } |z| < 2$$

$$\frac{1}{z-1} = \frac{1}{z} \cdot \frac{1}{1-\frac{1}{z}} = \left\{ (2^k) \right\} = \frac{1}{z} \cdot \sum_{n=0}^{\infty} \left(\frac{1}{z} \right)^n = \sum_{n=1}^{\infty} \frac{1}{z^n}, \text{ for } |z| > 1$$

$$f(z) = -\sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} - \sum_{n=1}^{\infty} \frac{1}{z^n}, \text{ for } 1 < |z| < 2$$

$$\begin{aligned} \frac{1}{1-c \cdot w} &= \sum_{n=0}^{\infty} (c \cdot w)^n, \text{ for } |w| < \frac{1}{|c|} \quad (1^*) \\ \frac{1}{1-\frac{b}{w}} &= \sum_{n=0}^{\infty} \left(\frac{b}{w} \right)^n, \text{ for } |w| > |b| \quad (2^*) \end{aligned}$$

note: $\frac{1}{(z-1)(z-2)} = \frac{1}{z-2} - \frac{1}{z-1}$
Partial fraction expansion

→ So our objective is to obtain the following formula format:

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n \cdot (z-z_0)^n + \sum_{n=1}^{+\infty} \frac{a_{-n}}{(z-z_0)^n}$$

• Things to note:

$$\bullet f(z) = \underbrace{\sum_{n=-\infty}^{+\infty} a_n \cdot (z-z_0)^n}_{\text{analytic part}} + \underbrace{\sum_{n=1}^{+\infty} \frac{a_{-n}}{(z-z_0)^n}}_{\text{principal part}}$$

$$\bullet f(z) = \sum_{\substack{n=-\infty \\ \text{starts at } \infty}}^{+\infty} a_n \cdot (z-z_0)^n + \sum_{\substack{n=1 \\ \text{starts at } 1}}^{+\infty} \frac{a_{-n}}{(z-z_0)^n}$$

